

New Method of Modelling Dissipative Hydrodynamics

T. Osada^a and G. Wilk^{b,1}

^a*Theoretical Physics Laboratory, Faculty of Knowledge Engineering, Musashi Institute of Technology,
Setagaya-ku, Tokyo 158-8557, Japan*

^b*The Andrzej Soltan Institute for Nuclear Studies, Hoża 69, 00681, Warsaw, Poland*

Abstract

We propose to model the dissipative hydrodynamics used in description of the multiparticle production processes (d -hydrodynamics) by a special kind of the perfect nonextensive fluid (q -fluid) where q denotes the nonextensivity parameter appearing in the nonextensive Tsallis statistics. The advantage of q -hydrodynamics lies in its formal simplicity in comparison to the d -hydrodynamics. We argue that parameter q describes summarily (at least to some extent) all dynamical effects behind the viscous behavior of the hadronic fluid.

Key words: nonextensive statistical mechanics, hydrodynamic models, high energy collisions, multiparticle production processes

PACS: 24.10.Nz, 25.75.-q

The ideal hydrodynamic model successfully reproduces most of the RHIC experimental data [1] suggesting that matter created there resembles a strongly interacting quark-gluon fluid [2] rather than a free parton gas. However, there are also hints that, after all, a hadronic fluid cannot be totally ideal [3] and should be described by some kind of dissipative hydrodynamic model (or d -hydrodynamics) [4]. A number of such models have been proposed recently [5,6]. Although very promising, they have many difficulties, both in what concerns their proper formulation and applications [6]. It turns out, however, that one can obtain constitutive equations of causal d -hydrodynamics starting from the relativistic perfect nonextensive hydrodynamics (q -hydrodynamics) supplemented by the nonextensive/dissipative correspondence (NexDC) proposed in [7] and taking place between the perfect q -hydrodynamics (based on the nonextensive Tsallis statistics [8] and described by the nonextensivity parameter q) and the usual d -hydrodynamics (based on the extensive Boltzmann-Gibbs statistics (BG) to which Tsallis statistics converges for $q \rightarrow 1$). .

Email addresses: `osada@ph.ns.musashi-tech.ac.jp` (T. Osada), `wilk@fuw.edu.pl` (G. Wilk).

¹ Presenting Author.

Our q -hydrodynamics is causal and preserves the simplicity of d -hydrodynamics facilitating its numerical applications [7]. It is based on the relativistic nonextensive kinetic theory proposed in [9], which replaces the usual notion of local thermal equilibrium by a kind of stationary state. This stationary state includes, by means of the parameter q [10], all kinds of possible strong intrinsic fluctuations and long-range correlations existing in a hadronizing system. In this approach the *perfect q -hydrodynamic equations* (i.e., equations for the perfect nonextensive q -fluid without any additional currents) are given by:

$$\mathcal{T}_{q;\mu}^{\mu\nu} = \left[\varepsilon_q(T_q) u_q^\mu u_q^\nu - P_q(T_q) \Delta_q^{\mu\nu} \right]_{;\mu} = 0, \quad (1)$$

where $\mathcal{T}_q^{\mu\nu}$ is q -version of the energy momentum tensor with $\varepsilon_q(T_q)$, $P_q(T_q)$, $T_q = T_q(x)$ and $u_q^\mu(x)$ being, respectively, the q -versions of the nonextensive energy density, pressure and temperature field [7] and an accompanying hydrodynamic q -flow four vector field (which allows decomposition (1) to be performed; $\Delta_q^{\mu\nu} \equiv g^{\mu\nu} - u_q^\mu u_q^\nu$). If $T(x)$ and $u^\mu(x)$ are the corresponding temperature and velocity fields in the case of BG statistics ($q = 1$) with ε and P being the usual energy density and pressure then, as proposed in [7], one can map a q -flow (with q -temperature T_q [7]) into some dissipative d -flow by requiring that the following relations hold:

$$P(T) = P_q(T_q), \quad \varepsilon(T) = \varepsilon_q(T_q) + 3\Pi, \quad (2)$$

where Π is to be regarded as the bulk pressure of d -hydrodynamics which is defined by

$$\Pi \equiv \frac{1}{3} w_q [\gamma^2 + 2\gamma]. \quad (3)$$

Here $\delta u_q^\mu \equiv u_q^\mu - u^\mu$ is the four velocity difference between the non-extensive parameter $q > 1$ and $q = 1$ flows, $\gamma(x) \equiv \delta u_q^\mu(x) u_\mu(x) = -\frac{1}{2} \delta u_{q\mu} \delta u_q^\mu$ and $w_q \equiv \varepsilon_q + P_q$ is the q -enthalpy.

Now, it can be shown [7] that, if $T(x)$ and $\gamma(x)$ satisfy relations (2) in the whole space-time region, one can always transform equation of perfect q -hydrodynamics Eq. (1) into the following equation of d -hydrodynamics [5]:

$$\left[\varepsilon(T) u^\mu u^\nu - [P(T) + \Pi] \Delta^{\mu\nu} + 2W^{(\mu} u^{\nu)} + \pi^{\mu\nu} \right]_{;\mu} = 0, \quad (4)$$

in which one recognizes the energy flow vector W^μ and the (symmetric and traceless) shear pressure tensor $\pi^{\mu\nu}$,

$$W^\mu = w_q [1 + \gamma] \Delta_\sigma^\mu \delta u_q^\sigma, \quad \pi^{\mu\nu} = w_q \delta u_q^{<\mu} \delta u_q^{\nu>} \quad (5)$$

(where $\Delta^{\mu\nu} \equiv g^{\mu\nu} - u^\mu u^\nu$, $A^{(\mu} B^{\nu)} \equiv \frac{1}{2} (A^\mu B^\nu + A^\nu B^\mu)$, and $a^{<\mu} b^{\nu>} \equiv [\frac{1}{2} (\Delta_\lambda^\mu \Delta_\sigma^\nu + \Delta_\sigma^\mu \Delta_\lambda^\nu) - \frac{1}{3} \Delta^{\mu\nu} \Delta_{\lambda\sigma}] a^\lambda b^\sigma$). The d -hydrodynamics represented by Eq. (4) can be therefore regarded as a viscous counterpart of the perfect q -hydrodynamics represented by Eq. (1). We call this relation the *nonextensive/dissipative correspondence*, (NexDC). With the bulk pressure Eq. (3) and NexDC relations Eq. (2) one gets the q -enthalpy $w_q = w/[1+\gamma]^2$ with $w \equiv Ts = \varepsilon + P$ being the usual enthalpy in BG statistics. Accordingly, the bulk pressure Eq. (3) can be written as

$$\Pi = w\Gamma, \quad \text{where} \quad \Gamma \equiv \frac{1}{3} \frac{\gamma(\gamma+2)}{(\gamma+1)^2}. \quad (6)$$

Finally, the NexDC leads to the following relations between components of the dissipative tensor:

$$W^\mu W_\mu = -3\Pi w, \quad \pi^{\mu\nu} W_\nu = -2\Pi W^\mu, \quad \pi_{\mu\nu} \pi^{\mu\nu} = 6\Pi^2. \quad (7)$$

Referring to [7] for details and examples of numerical applications, we would like to close with two remarks. First, albeit in our q -hydrodynamic the q -fluid is ideal (there is no production of q -entropy), the usual entropy in the corresponding d -fluid is produced (as result, the total multiplicity increases with $(q - 1)$). Second, this approach results in a very simple relation between the ratios of shear and bulk viscosities over the entropy density, η/s and ξ/s , respectively:

$$\frac{1}{\xi/s} + \frac{3}{\eta/s} = \frac{w\sigma_{\text{full};\mu}^\mu}{\Pi^2}. \quad (8)$$

where σ_{full}^μ is the full order dissipative entropy current. Further investigations of q -hydrodynamics and its implications are under way.

Acknowledgment

GW would like to express his gratitude towards the organizers of the PANIC2008 conference (Eilat, Israel, 10-14 November 2008) where this work was presented. Partial support (GW) from the Ministry of Science and Higher Education under contract 1P03B02230 is also acknowledged.

References

- [1] T. Hirano, N. Kolk and A. Bilandzic, airXiv:0808.2684, to appear in Springer Lecture Notes in Physics.
- [2] T. D. Lee, Nucl. Phys. **A750**, 1 (2005).
- [3] T. Hirano and M. Gyulassy, Nucl. Phys. A **769**, 71 (2006).
- [4] C. Eckart, Phys.Rev. **58**, 919 (1940); W. Israel, Ann. Phys. (N.Y.) **100**, 310 (1976); J. M. Stewart, Proc. R. Soc. London, Ser. A **357**, 59 (1977); W. Israel and J. M. Stewart, *Ann Phys. (N.Y.)* **118**, 341 (1979).
- [5] A. Muronga, Phys. Rev. **C 69**, 034903 (2004) and **C 76**, 014910 (2007); U. Heinz, H. Song and A.K. Chaudhuri, Phys. Rev. **C 73**, 034904 (2006); H. Song and U. Heinz, Phys. Lett. B **658**, 279 (2008); R. Baier and R. Romatschke and U. A. Wiedemann, Phys. Rev. **C 73**, 064903 (2006); A. K. Chaudhuri, Phys. Rev. **C 74**, 044904 (2006).
- [6] W. A. Hiscock and L. Lindblom, Phys. Rev. **D 31**, 725 (1985); **D 35**, 3723 (1987); T. Koide, G. S. Denicol, Ph. Mota and T. Kodama, Phys. Rev. **C 75**, 034909 (2007); G.S.Denicol, T. Kodama, T. Koide and Ph. Mota, Phys. Rev. **C 78**, 034901 (2008). P.Ván and T.S.Biro, Eur. Phys.J. Special Topics **155**, 201 (2008); T.S. Biro, E. Molnar and P. Ván, Phys. Rev. **C 78**, 014909. (2008)
- [7] T. Osada and G. Wilk, Phys. Rev. **C 77**, 044903 (2008) and Prog. Theor. Phys. Suppl. **174**, 168 (2008); cf. also: arXiv:0805.2253[nucl-phys] (to be published in Ind. J. Phys.), arXiv:0810.3089[hep-ph] (to be published in Centr. Eur. J. Phys.); arXiv:0810.5192[nucl-th].
- [8] C. Tsallis, J. Stat. Phys. **52** 479 (1988). For updated bibliography on this subject see <http://tsallis.cat.cbpf.br/biblio.htm>.
- [9] A. Lavagno, Phys. Lett. **A301**, 13 (2002).

- [10] T. Kodama, H. T. Elze, C. E. Aguiar and T. Koide, Europhys. Lett. **70**, 439 (2005); T.S. Biró and G. Purcsel, Phys. Rev. Lett. **95** 162302 (2005); Phys. Lett. A **372**, 1174 (2008); T. Sherman and J. Rafelski, Lecture Notes in Physics **633** (2004) 377.